Available online at www.sciencedirect.com



Journal of Acupuncture and Meridian Studies

journal homepage: www.jams-kpi.com



Research Article

Effects of A Short-term Cardio Tai Chi Program on Cardiorespiratory Fitness and Hemodynamic Parameters in Sedentary Adults: A Pilot Study



Timothy Sam-Kit Tin¹, Chi-Hsiu Daniel Weng^{1,2}, Patricia dos Santos Vigário³, Arthur de Sá Ferreira^{3,4,*}

¹ University of East-West Medicine, Sunnyvale, CA, 94085, USA

² Department of Kinesiology, San Jose State University, CA, 95112, USA

³ Postgraduate Program in Rehabilitation Sciences, Augusto Motta University Center, Praça

das Nações 34, Bonsucesso, Rio de Janeiro, RJ, 21041-010, Brazil

⁴ Salgado de Oliveira University, Rua Marechal Deodoro 263, Centro, Niterói, RJ, 24030-060, Brazil

Available online 18 December 2019

Received: Dec 6, 2018 Revised: Jul 20, 2019 Accepted: Dec 11, 2019

KEYWORDS

complementary therapies; exercise; mind-body therapies; rehabilitation

Abstract

This study evaluates the effects of a short-term Cardio *Tai Chi* program on the cardiorespiratory fitness and hemodynamic parameters in sedentary adults. Thirty-one sedentary participants (age: 58 ± 9 years, body mass: 63 ± 12 kg) were subjected to an exercise program during 10 sessions over a 10-day period within 2 weeks. The Cardio *Tai Chi* program consisted in a series of three to five intervals lasting 90 s each at ~70% maximal heart rate separated by 2-min of low-intensity recovery. Primary outcome measures were cardiorespiratory fitness (peak oxygen uptake, VO_{2peak}) assessed by the Rockport walking test and resting hemodynamic parameters (systolic, diastolic, mean, and pulse pressures). We observed a significant difference of means on post-pre VO_{2peak} [4.5 ml/kg/ min, 95% confidence interval (CI): 3.1 to 5.8, p = 0.004], systolic blood pressure (-5.5 mmHg, 95% CI:-7.3 to -3.8, p = 0.010) and pulse pressure (-3.7 mmHg, 95% CI:

* Corresponding author. Program in Rehabilitation Sciences, Augusto Motta University Center, Praça das Nações 34, Bonsucesso, Rio de Janeiro, RJ, 21041-010, Brazil.

E-mail: arthur_sf@icloud.com (A.S. Ferreira).

pISSN 2005-2901 eISSN 2093-8152

https://doi.org/10.1016/j.jams.2019.12.002

© 2020 Medical Association of Pharmacopuncture Institute, Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

-5.2 to -2.3, p = 0.028). No significant differences were observed for diastolic pressure (-1.8 mmHg, 95% Cl: -2.6 to -1.0, p = 0.226), mean blood pressure (2.5 mmHg, 95% Cl: 1.4 to 3.6, p = 0.302), or resting heart rate (-0.9 beat/min, 95% Cl: -2.0 to 0.1, p = 0.631). Our findings suggest that engaging in a short-term Cardio *Tai Chi* program can improve cardiorespiratory fitness and hemodynamic parameters in sedentary adults.

1. Introduction

Physical inactivity-an activity level insufficient to meet current recommendations- is considered the major public health problem of this century [1], accounting for 3.2% to 7.8% of coronary heart disease worldwide [2]. There is compelling evidence that both physical activity and exercise are associated to numerous positive health-related biopsychosocial outcomes such as lowering blood pressure and cardiovascular disease risk factors, improving the cardiorespiratory fitness, preserving bone mass, and reducing the risk of falling, improving anxiety, guality of life, and delaying allcause mortality [3]. A report on the pandemic of physical inactivity strongly urges the academics to provide supporting evidence for effective programs to advance global health through physical activity [4]. Because exercise can be prescribed in the treatment of many diseases, including psychiatric, neurological, metabolic, musculoskeletal, cancer, pulmonary, and cardiovascular [5], there is a need for programs that can promote the regular practice of physical activity in the general population and particularly those at a high risk of cardiovascular diseases.

Tai Chi Chuan-or Tai Chi-is a Chinese mind-body practice enrolling meditation, static and dynamic body postures associated with controlled breathing that is most closely related to exercise [6,7]. The practice of *Tai Chi* comprises sequences of stances, *i.e.*, coordinated upper- and lowerlimb movements in predominantly slow and smooth trajectories [8-11]. A large multicenter study from 13 countries summing up 6410 participants performing either Tai Chi or Qi Gong reported health benefits related to bone density, cardiorespiratory effects, physical function, falls prevention, postural balance, and related risk factors, quality of life, self-efficacy, patient-reported outcomes, psychological symptoms, and immune- and inflammationrelated responses [12]. As related to cardiovascular diseases, recent systematic reviews and meta-analyses show that Tai Chi practice can improve systolic blood pressure (SBP) and diastolic blood pressure (DBP) and cardiorespiratory fitness [13-15]. Although complementary and alternative medicine therapies are promising for patients with cardiac rehabilitation, it still lacks high-quality evidence in the study of the relationship between Tai Chi and cardiorespiratory fitness [14-17].

Improving cardiorespiratory fitness and reducing other cardiovascular risk factors such as blood pressure are both strongly associated to a regular engaging in physical activity with defined frequency, intensity, time, type, volume, pattern, and progression of cardiorespiratory exercise [3] All the aforementioned requirements are met by traditional *Tai Chi* practice except for the exercise intensity [18,19]. We thus speculate that *Tai Chi* practice might elicit the minimal exercise intensity level (*i.e.*, threshold), provided the traditional stances are performed under more vigorous movements thus promoting cardiorespiratory fitness [3]. As

a corollary, risk factors related to cardiovascular diseases might be modified as well. Therefore, this pilot study aims to evaluate the effects of a short-term Cardio *Tai Chi* program on the cardiorespiratory fitness and hemodynamic parameters in sedentary adults. We hypothesized that the regular practice of *Tai Chi* at aerobic target intensity levels might elicit physiological responses that positively affect cardiorespiratory fitness and hemodynamic parameters, both associated with reduced cardiovascular risk.

2. Materials and Methods

2.1. Ethical approval

The Institutional Review Board at the University of East-West Medicine has approved this project before its execution in accordance with the ethical standards set in the Helsinki Declaration of 1975. Participants provided written informed consent before being admitted to this study.

2.2. Study design and sample size estimates

This pilot study is an uncontrolled, before-and-after study design (quasi-experimental) [20,21] in which a shortterm program of exercises was applied in 10 sessions over a 10-day period within 2 weeks (Fig. 1). Measurements were obtained from baseline testing and after the 10 sessions were finished. Three assessors performed the measurements independently, and they were not aware of the purpose of the measurements. Participants were not informed about their testing results during their enrollment.

The sample size was estimated for the main outcome of this study, i.e., maximal oxygen uptake in metabolic equivalent (MET) corresponding to 3.5 mL/kg/min in a healthy 40-year-old 70kg man [22]. A minimal sample size of 31 participants in this single-arm, before-and-after study was obtained from $\beta = 0.20$, $\alpha_{two-tailed} = 0.05$, a small effect size = 0.2 as in a previous study [23] and S(Δ) = 0.4 using an online calculator (http://www.sample-size.net/ sample-size-study-paired-t-test/).

2.3. Participants

Thirty-one participants (age: 58 ± 9 years, body mass: 63 ± 12 kg) were enrolled in this study, and all participants completed the study protocol without missing sessions. Participants were contacted from posters, churches, and word of mouth. Eligibility criteria comprised ages between 30 and 80 years, no diagnosis of cardiovascular or metabolic diseases, no use of medication regularly, absence of musculoskeletal pain or condition that limit exercise practice, commitment of attending the *Tai Chi* sessions, and a



Figure 1 Study flowchart.

self-reported sedentary lifestyle (*i.e.*, not engaging in physical activity for the last 6 months).

2.4. Intervention: the Cardio *Tai Chi* exercise program

The Cardio *Tai Chi* program was designed after the integration of traditional *Tai Chi* practice [24] and features of aerobic exercise. The class of Cardio *Tai Chi* comprises selected *Tai Chi* stances performed at a fast pace for a higher intensity exercise (Fig. 2). Exercise intensity is moderated by varying the number of intervals of fast pace (two *Tai Chi* movements) that are interleaved by slow pace movements (one *Tai Chi* movement), with transitional motion (two *Tai Chi* movements). The high-intensity movement of each stance is progressively moving the body from moderate speed to higher speed within 90 s in each section.

The training program consisted in 10 sessions over a 10day period within 2 weeks. Participants were familiarized with the *Tai Chi* stances before the first session. Sessions were performed daily from Saturday to Tuesday and Thursday (with Wednesdays and Fridays as rest days). The intensity of the sessions was distributed as follows (Table 1): three repeated sections in Level #1 (3 intervals of fast pace movements lasting 90 s); four sections in Level #2 (4 intervals of fast pace movements lasting 90 s) and five sections in Level #3 (5 intervals of fast pace movements lasting 90 s). Within each interval, the exercise intensity was increased until eliciting the \sim 70% maximal heart rate (HR) as measured during the assessment (section 2.5). Each interval of fast pace movements was separated by 2 minutes of low-intensity recovery. Participants had a 30-s resting period and 30 s to switch the *Tai Chi* stance into slow motion before and after the rest. There is a 30-s transition time between fast and slow pace. All sessions started and ended with a 5-minute period of warm-up and cooldown using the *Tai Chi* stances. The total time of exercise in Levels 1-3 is ~20, 24, and ~27 min, respectively.

2.5. Assessments

The primary outcome consisted of estimated VO_{2peak} as determined from a continuous submaximal exercise test on an electronically treadmill (Pro-Form 505 CST) using the Rockport walking test [25,26]. The test consisted in walking as quickly as possible for 1600 m (running or jogging was not permitted) wearing the Wireless Blood Pressure Monitor BP5 (iHealth Labs Inc., CA, USA). The time spent to complete the target distance and the HR achieved immediately after the test were both recorded and then computed in an agespecific regression equation allowing for the estimation of \dot{VO}_{2peak} . The \dot{VO}_{2peak} was expressed in mL/kg/min and by calculating the MET [3]. Differences between values higher than 1 MET before and after the program were considered clinically relevant [27].

Secondary outcomes comprised resting HR and resting blood pressure. During the supervised training intervention, HR was recorded using Deluxe Pulse Oximeter (Veridian Deluxe Pulse Oximeter #11-50D) to ensure that participants were exercising at the prescribed target intensity. SBP and DBP were also assessed using the Wireless Blood Pressure



Figure 2 Slow, fast, and transitional stances used in Cardio *Tai Chi* Program. Exercise intensity is moderated by varying the number of intervals of fast pace (two *Tai Chi* movements) that are interleaved by slow pace movements (one *Tai Chi* movement), with transitional motion (two *Tai Chi* movements). The high-intensity movement of each stance is progressively moving the body from moderate speed to higher speed within 90 s in each section.

Monitor BP5 (iHealth Labs Inc., CA, USA). Pulse pressure (PP = SBP - DBP) and mean blood pressure (MBP = DBP + PP/3) were also calculated [28].

2.6. Statistical analysis

Data were typed in an electronic worksheet for statistical analysis (Excel, Microsoft Corp., USA). Data are summarized as mean \pm SD or frequency (%) depending on the variable type. Difference in Means (MD) and standardized mean difference (SMD) with 95% confidence intervals (95% CI) were also calculated as effect size estimates. Beforeand-after comparisons were evaluated using the *t* test for dependent samples (two-tailed analysis). Pearson's coefficient was used to quantify the correlation between values of the outcome variables before and after the program. Statistical significance was set to p < 0.05. Substantive evidence was assessed by effect sizes (SMD) and is interpreted as per the suggested values [23].

3. Results

Table 2 summarizes the descriptive and comparative analyses of cardiorespiratory fitness and hemodynamic parameters. A significant MD was observed on post-pre \dot{VO}_{2peak} (MD 4.5 ml/kg/min, 95% CI: 3.1 to 5.8, p = 0.004), with a MD higher than the 1-MET minimum clinically important difference (MD: 1.3 MET, 95% CI: -0.1 to 2.6, p = 0.004).

Significant differences were also observed for SBP (MD: -5.5 mmHg, 95% CI: -7.3 to -3.8, p = 0.010) and pulse pressure (MD: -3.7 mmHg, 95% CI: -5.2 to -2.3, p = 0.028), all with large effect sizes (SMD: 0.94 to 1.27). No significant differences were observed for diastolic pressure (MD: -1.8 mmHg, 95% CI: -2.6 to -1.0, p = 0.226) and mean blood pressure (MD: 2.5 mmHg, 95% CI: 1.4 to 3.6, p = 0.302), both showing a medium effect size (SMD: 0.60 to 0.73). Resting HR was also not statistically different on post-pre analysis (MD: 0.9 beat/min, 95% CI: -2.0 to 0.1, p = 0.631), with a small effect size (SMD: 0.27).

Correlation analysis between values before and after the program showed moderate to strong correlation coefficients for \dot{VO}_{2peak} (r = 0.857), SBP (r = 0.830), pulse pressure (r = 0.839), HR (r = 0.600), DBP (r = 0.599), and mean blood pressure (r = 0.352).

4. Discussion

This pilot study evaluated the effects of a short-term Cardio *Tai Chi* program on the cardiorespiratory fitness and hemodynamic parameters in sedentary adults. We found both statistical and clinical evidence favoring the hypotheses that the short-term practice of *Tai Chi* at minimum aerobic target intensity levels elicits physiological responses that positively affects cardiorespiratory fitness and hemodynamic parameters. To the best of our knowledge, this is the first study to propose the *Tai Chi* practice under aerobic target intensity levels and to evaluate its effect on cardiorespiratory fitness and hemodynamic parameters, both associated to cardiovascular risk. This study fills an important gap in literature [13,14] regarding whether the already acknowledged biopsychosocial benefits of practicing traditional *Tai Chi* are observed in other circumstances of practice.

We observed a substantive increase in estimated \dot{VO}_{2peak} that was also larger than the minimum clinically important difference of 1 MET (each 1-MET increase in exercise capacity is related to a 10 to 20% improvement in survival from cardiovascular diseases) [27]. This finding is slightly

Table 1 Cardio *Tai Chi* program schedule. All sessions started and ended with a 5-minute period of warm-up and cooldown using the *Tai Chi* stances. The intensity of the sessions was distributed as follows: three repeated sections in Level #1 (3 intervals of fast pace movements lasting 90 s), four sections in Level #2 (4 intervals of fast pace movements lasting 90 s), and five sections in Level #3 (5 intervals of fast pace movements lasting 90 s).

Task	Intensity	Tai Chi stances					
duration (mm:ss)		Level 1 (Classes 1 to 3)	Level 2 (Classes 4 to 6)	Level 3 (Classes 7 to 10)			
05:00	Warm-up	Stretching exercises	Stretching exercises	Stretching exercises			
00:30	Moderate	Cross hand	Cross hand	Cross hand			
00:30	High intensity	Jumping rooster	Playing fiddle	Catching bird			
00:30	Moderate	Jumping rooster	Side kicking	White crane			
00:30	Slow	Jumping	Playing	Catching			
	down	rooster/	fiddle/	bird/			
		mapping	mapping	mapping			
		mirror	mirror	mirror			
00:30	Slow	Cloud	Open	Embrace			
	motion	hand	window	tiger			
00:30	Break	Rest	Rest	Rest			
00:30	Slow	Cloud	Open	Embrace			
	motion	hand	window	tiger			
Repetitions	-	3	3	3			
·		repetitions	repetitions,	repetitions,			
			Level 1	Level 1 and 2			
05:00	Cool	Deep	Deep	Deep			
	down	breathing exercise	breathing exercise	breathing exercise			
Total time	-	20:30	24:00	27:30			

better than those reported after 16-week high-intensity interval training program or continuous moderateintensity exercise training in adults with hypertension [29]. This suggests that the Cardio Tai Chi program using several selected stances from the simplified form of Tai Chi at aerobic target intensity levels was capable of promoting cardiorespiratory fitness. In the traditional practice of Tai *Chi*, the upper and lower extremities perform movements, whereas the body moves resembling walking at a speed that is relatively slower compared with activities of daily living [8-11]. Because walking speed is strongly correlated to both aerobic capacity and mitochondrial activity [30], we speculate that movement speed in the Cardio Tai Chi played a major role in the observed cardiorespiratory effects. Interestingly, studies comparing the effects of traditional Tai Chi practice versus brisk walking on cardiorespiratory fitness found no difference in elderlies [31,32].

The observed reduction in systolic and pulse pressures was also substantive and similar to those reported by systematic reviews and meta-analyses [13-15] on the traditional Tai Chi practice. Systolic and pulse pressures are well known, strong, independent predictors of cardiovascular morbidity [33]. The finding that Cardio *Tai Chi* contributes to lowering blood pressure values strongly suggest this program as an option for controlling cardiac risk factors by engaging in a regular physical activity. Nonetheless, diastolic and mean blood pressures did not improve after this 2-week Cardio Tai Chi program, as well as resting heart rate. The traditional practice of Tai Chi for 5 min acute and temporarily increases the heart rate variability (HRV) in young and older adults [34], which is in agreement that larger HRV during exercise is associated with better physical fitness [35]. We thus speculate whether a longer trial could elicit stronger substantive effects on those cardiac risk factors if any.

The findings presented herein should be interpreted in light of the factors jeopardizing the internal validity of uncontrolled before-and-after study design [21]. Therefore, we cannot rule out history, maturation, and instru*ment decay* as rival hypotheses, but owing to the relative short-term between assessments all were considered implausible. Similarly, bias due to testing and reactivity were both also considered unlikely owing to the blinding of the participants to their own results. However, statistical regression (toward the mean) is weakly plausible because of the strong correlation between values of all studied variables before and after the program, mainly mean blood pressure. Nonetheless, our results strongly encourage conducting experimental trials with larger samples, direct methods for VO_{2peak} measurement, longer follow-up periods (e.g., 4, 8 or 12 weeks) and control group to further explore the effects of Cardio Tai Chi on cardiorespiratory fitness and other cardiovascular risk factors and to observe whether the observed effects are transient or long lasting. The integration of the aerobic exercise and the traditional Tai Chi is a promising contribution to sport and rehabilitation fields.

In summary, the short-term practice of *Tai Chi* at minimum aerobic target intensity levels elicits physiological responses that positively affects cardiorespiratory fitness and hemodynamic parameters, all associated with reduced cardiovascular risk in sedentary adults.

Table 2Descriptive and comparative analysis.

Variable	Study phase		P-value	Standardized	Effect size estimate
	Before	After		mean difference (SMD)	
Cardiorespiratory fitness					
<i>VO_{2peak}</i> , ml/kg/min	$\textbf{37} \pm \textbf{14}$	41 ± 15	0.004*	1.16	Large
VO _{2peak} , MET	$\textbf{10.4} \pm \textbf{4.1}$	$\textbf{11.7} \pm \textbf{4.3}$	0.004*	0.33	Small
Hemodynamic parameters at rest					
Systolic blood pressure, mmHg	133 \pm 19	127 ± 19	0.010*	-1.27	Large
Diastolic blood pressure, mmHg	77 ± 9	75 ± 9	0.226	-0.60	Medium
Pulse pressure, mmHg	56 \pm 16	52 ± 15	0.028*	-0.94	Large
Mean blood pressure, mmHg	91 ± 12	93 ± 11	0.302	0.73	Medium
Heart rate, beat/min	77 ± 13	76 ± 10	0.631	-0.27	Small
Data are presented as mean \pm SD					

Data are presented as mean \pm SD.

* Statistical significance at p < 0.05.

MET, metabolic equivalent.

Conflict of interest

We declare that we have no conflicts of interest regarding this study.

Acknowledgments

This study was supported by the Fundação Carlos Chagas Filho de Apoio à Pesquisa do Estado do Rio de Janeiro (grant number E-26/202.769/2015). The funding source had no role in the study design, in the collection, analysis and interpretation of data, in the writing of the manuscript, and in the decision to submit the manuscript for publication.

References

- [1] Trost SG, Blair SN, Khan KM. Physical inactivity remains the greatest public health problem of the 21st century: Evidence, improved methods and solutions using the "7 investments that work" as a framework. Br J Sports Med 2014;48:169–70. https: //doi.org/10.1136/bjsports-2013-093372.
- [2] Lee I-M, Shiroma EJ, Lobelo F, Puska P, Blair SN, Katzmarzyk PT. Impact of Physical Inactivity on the World's Major Non-Communicable Diseases. Lancet 2012;380:219–29. https://doi.org/10.1016/S0140-6736(12)61031-9. Impact.
- [3] Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee IM, et al. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: Guidance for prescribing exercise. Med Sci Sports Exerc 2011; 43:1334–59. https: //doi.org/10.1249/MSS.0b013e318213fefb.

//doi.org/10.1249/MSS.000136318213161D.

- [4] Kohl HW, Craig CL, Lambert EV, Inoue S, Alkandari JR, Leetongin G, et al. The pandemic of physical inactivity: global action for public health. Lancet 2012;380:294–305. https: //doi.org/10.1016/S0140-6736(12)60898-8.
- [5] Pedersen BK, Saltin B. Exercise as medicine Evidence for prescribing exercise as therapy in 26 different chronic diseases. Scand J Med Sci Sport 2015;25:1–72. https: //doi.org/10.1111/sms.12581.

- [6] de Sá Ferreira A. Evidence-based practice of Chinese medicine in physical rehabilitation science. Chin J Integr Med 2013;19: 723-9. https://doi.org/10.1007/s11655-013-1451-5.
- [7] Cheng TO. Tai Chi: The Chinese ancient wisdom of an ideal exercise for cardiac patients. Int J Cardiol 2007;117:293–5. https://doi.org/10.1016/j.ijcard.2006.06.021.
- [8] Wu G, Ren X. Speed effect of selected Tai Chi Chuan movement on leg muscle activity in young and old practitioners. Clin Biomech 2009;24:415–21. https: //doi.org/10.1016/j.clinbiomech.2009.03.001.
- [9] Wolf S, Coogler C, Xu T. Exploring the basis for Tai Chi Chuan as a terapeutic exercise approach. Arch Phys Med Rehabil 1997;78:886–92.
- [10] Wu G, Hitt J. Ground contact characteristics of Tai Chi gait. Gait Posture 2005;22:32–9. https: //doi.org/10.1016/j.gaitpost.2004.06.005.
- [11] Wu G, Liu W, Hitt J, Millon D. Spatial, temporal and muscle action patterns of Tai Chi gait. J Electromyogr Kinesiol 2004; 14:343–54. https://doi.org/10.1016/j.jelekin.2003.09.002.
- [12] Jahnke R, Larkey L, Rogers C, Etnier J, Lin F. A comprehensive review of health benefits of qigong and tai chi. Am J Health Promot 2010;24. https://doi.org/10.4278/ajhp.081013-LIT-248.
- [13] Yeh GY, Wang C, Wayne PM, Phillips RS. The effect of Tai Chi exercise on blood pressure: A systematic review. Prev Cardiol 2008;11:82–9. https://doi.org/10.1111/j.1751-7141.2008.07565.x.
- [14] Zheng G, Li S, Huang M, Liu F, Tao J, Chen L. The effect of Tai Chi training on cardiorespiratory fitness in healthy adults: A systematic review and meta-analysis. PLoS One 2015;10:1–20. https://doi.org/10.1371/journal.pone.0117360.
- [15] Yang Y, Wang Y, Wang S, Shi P, Wang C. The Effect of Tai Chi on Cardiorespiratory Fitness for Coronary Disease Rehabilitation: A Systematic Review and Meta-Analysis. Front Physiol 2018;8: 1–10. https://doi.org/10.3389/fphys.2017.01091.
- [16] Arthur HM, Patterson C, Stone JA. The role of complementary and alternative therapies in cardiac rehabilitation: a systematic evaluation. EurJ Cardiovasc PrevRehabil 2006;13: 3–9. https://doi.org/10.1097/01.hjr.0000198917.67987.f8.
- [17] Wang J, Xiong X. Evidence-based chinese medicine for hypertension. Evid Based Complement Alternat Med 2013;2013. https://doi.org/10.1155/2013/978398. 978398.
- [18] Hong Y, Li JX, Robinson PD. Balance control, flexibility, and cardiorespiratory fitness among older Tai Chi practitioners. Br J Sports Med 2000;34:29–34.
- [19] Lan C, Lai JS, Wong MK, Yu ML. Cardiorespiratory function, flexibility, and body composition among geriatric Tai Chi

Chuan practitioners. Arch Phys Med Rehabil 1996;77:612-6. https://doi.org/10.1016/S0003-9993(96)90305-6.

- [20] Arain M, Campbell MJ, Cooper CL, Lancaster GA. What is a pilot or feasibility study? A review of current practice and editorial policy. BMC Med Res Methodol 2010;10:67. https: //doi.org/10.1186/1471-2288-10-67.
- [21] Campbell DT, Stanley JC. Experimental and Quasi-Experimental Design for Research. 1967. https: //doi.org/10.1037/022808.
- [22] Byrne NM, Hills AP, Hunter GR, Weinsier RL, Schutz Y. Metabolic equivalent: one size does not fit all. J Appl Physiol 2005; 99:1112–9. https: //doi.org/10.1152/japplphysiol.00023.2004.
- [23] Faraone SV. Interpreting estimates of treatment effects: implications for managed care. Pharm Ther 2008;33:700-11.
- [24] Weng CD. The Official Cardio Tai- Chi Handbook. 1st ed. San Jose: Hunter Textbooks Inc.; 2007.
- [25] Kline GM, Porcari JP, Hintermeister R, Freedson PS, Ward A, McCarron RF, et al. Estimation of VO2max from a one-mile track walk, gender, age, and body weight. Med Sci Sports Exerc 1987;19:253–9. PMID: 3600239.
- [26] Pober DM, Freedson PS, Kline GM, McInnis KJ, Rippe JM. Development and validation of a one-mile treadmill walk test to predict peak oxygen uptake in healthy adults ages 40 to 79 years. Can J Appl Physiol 2002;27:575–89. https: //doi.org/10.1139/h02-033.
- [27] Artero EG, Jackson AS, Sui X, Lee D, O'Connor DP, Lavie CJ, et al. Longitudinal Algorithms to Estimate Cardiorespiratory Fitness: Associations with Nonfatal Cardiovascular Disease and Disease- Specific Mortality. J Am Coll Cardiol 2014;63:2289–96. https://doi.org/10.1016/j.jacc.2014.03.008.Longitudinal.

- [28] Salvi P. Mean Arterial Pressure. Pulse Waves 2012:3-8. https: //doi.org/10.1007/978-88-470-2439-7.
- [29] Ciolac EG. High-intensity interval training and hypertension: maximizing the benefits of exercise? Am J Cardiovasc Dis 2012; 2:102-10. www.AJCD.us/ISSN:2160-200X/AJCD1202003.
- [30] Coen PM, Jubrias SA, Distefano G, Amati F, Mackey DC, Glynn NW, et al. Skeletal Muscle Mitochondrial Energetics Are Associated With Maximal Aerobic Capacity and Walking Speed in Older Adults. Journals Gerontol Ser A 2013;68:447–55. https://doi.org/10.1093/gerona/gls196.
- [31] Audette JF, Jin YS, Newcomer R, Stein L, Duncan G, Frontera WR. Tai Chi versus brisk walking in elderly women. Age Ageing 2006; 35:388–93. https://doi.org/10.1093/ageing/afl006.
- [32] Ji Z, Li A, Feng T, Liu X, You Y, Meng F, et al. The benefits of Tai Chi and brisk walking for cognitive function and fitness in older adults. PeerJ 2017;5. https://doi.org/10.7717/peerj.3943. e3943.
- [33] Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL, et al. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: The JNC 7 Report. J Am Med Assoc 2003; 289:2560–72. https://doi.org/10.1001/jama.289.19.2560.
- [34] Vaananen J, Xusheng S, Wang S, Laitinen T, Pekkarinen H, Lansimies E, et al. Taichiquan acutely increases heart rate variability. Clin Physiol Funct Imaging 2002;22:2–3. https: //doi.org/10.1046/j.1475-097X.2002.00355.x.
- [35] Tulppo MP, Mäkikallio TH, Seppänen T, Laukkanen RT, Huikuri HV, Coote JH, et al. Vagal modulation of heart rate during exercise: effects of age and physical fitness and training load. Am J Physiol Heart Circ Physiol 2012;274: H424-9.